

ownership interest, scientific Advisor or membership, and speakers bureau with Natera; and other interests/relationships with HossMed, Inc.

## FUNDING

None.

## REFERENCES

1. Bunnapradist S, Homkraisas P, Ahmed E, Fehringer G, Billings P, Tabriziani H: Using both the fraction and quantity of donor-derived cell-free DNA to detect kidney allograft rejection. *J Am Soc Nephrol* 32: 2439–2441, 2021
2. Bloom RD, Bromberg JS, Poggio ED, Bunnapradist S, Langone AJ, Sood P, et al.; Circulating Donor-Derived Cell-Free DNA in Blood for Diagnosing Active Rejection in Kidney Transplant Recipients (DART) Study Investigators: Cell-free DNA and active rejection in kidney allografts. *J Am Soc Nephrol* 28: 2221–2232, 2017
3. Sigdel TK, Archila FA, Constantin T, Prins SA, Liberto J, Damm I, et al.: Optimizing detection of kidney transplant injury by assessment of donor-derived cell-free DNA via massively multiplex PCR. *J Clin Med* 8: 19, 2018

See related letter to the editor, “Need for a Validation Study before Using the Two-Step Algorithm for dd-cfDNA to Screen for Acute Rejection,” on pages 2972–2973 and research letter, “Using both the Fraction and Quantity of Donor-Derived Cell-Free DNA to Detect Kidney Allograft Rejection,” in Vol. 32, Iss. 10, on pages 2439–2441.

Suphamai Bunnapradist<sup>1</sup> and Hossein Tabriziani<sup>2</sup>

<sup>1</sup>Division of Nephrology, Department of Medicine, David Geffen School of Medicine at University of California, Los Angeles at Los Angeles at California

<sup>2</sup>Medical Affairs, Natera, Inc., San Carlos, California

JASN 2973–2974, 2021.

doi: <https://doi.org/10.1681/ASN.2021081039>

# Incidence of Arteritis and Peritubular Capillaritis in ANCA-associated Vasculitis

ANCA-associated vasculitis (AAV) is a small-vessel vasculitis affecting multiple organ systems, including the kidney. Small vessels in the kidney include small arteries (interlobular artery, afferent and efferent arteriole), capillaries (glomerular and peritubular), and venules. Although crescentic ANCA GN is a common histologic finding reflecting glomerular small-vessel vasculitis, it is reasonable the manifestation of AAV could also contribute to interstitial small-vessel vasculitis. With great interest, we read the recent study by Boudhabhay *et al.*<sup>1</sup> reporting that arteritis

represents a subtype of AAV, affecting 13.5% of patients. In addition, implementation of arteritis significantly improved the ANCA renal risk score (ARRS) for the prediction of ESKD.<sup>1,2</sup> These important observations improve our understanding of mechanisms contributing to renal injury in AAV. Besides arteritis, vasculitis manifestation to peritubular capillaries (peritubular capillaritis) has also been reported.<sup>3</sup> In this study, we share our own experience of the incidence of arteritis and peritubular capillaritis in kidney biopsies of confirmed AAV. In kidney biopsies with the presence of both arteries and peritubular capillaries, incidence of arteritis was 11 out of 42 patients (26.2%) and peritubular capillaritis four out of 42 (9.5%), respectively. There was an overlap between interstitial arteritis and peritubular capillaritis in most patients, both detectable in three out of 42 (7.1%). Peritubular capillaritis without arteritis was only present in one out of 42 (2.4%) patients. In summary, in this study we confirm that arteritis reflects the predominant interstitial vasculitis manifestation in a considerable subset of patients with AAV.

For almost two decades, constant effort has been made to stratify ESKD risk in AAV. Histopathological subgrouping into four classes (focal, crescentic, mixed, and sclerotic) as defined by Berden *et al.*<sup>4</sup> was proposed to predict long-term renal survival rates. Unlike Berden’s classification, Brix *et al.*<sup>2</sup> suggested the ARRS by incorporation of baseline GFR to the histopathological findings (percentage of normal glomeruli, tubular atrophy/interstitial fibrosis) to predict ESKD in AAV. The recent study by Boudhabhay *et al.* expands our knowledge of renal interstitial lesions in AAV, revealing arteritis as an important factor associated with severe kidney injury in a considerable subset of AAV. Furthermore, the fact that implementation of arteritis significantly improved the ARRS implicates a contribution of arteritis to kidney injury independent of established scoring systems.<sup>2</sup> Although the long-term renal outcome is of relevance, systematic analysis of interstitial vasculitis manifestation with arteritis and peritubular capillaritis in association with glomerular among other tubulointerstitial lesions would also be of great interest.

## ETHICS APPROVAL AND CONSENT TO PARTICIPATE

The study was performed in accordance with Good Clinical Practice and Declaration of Helsinki principles for ethical research. This retrospective study was approved by the ethics committee of the University Medical Center Göttingen, Germany (no. 4/8/19).

## DISCLOSURES

All authors have nothing to disclose.

Published online ahead of print. Publication date available at [www.jasn.org](http://www.jasn.org).

**Correspondence:** Dr. Björn Tampe, Department of Nephrology and Rheumatology, University Medical Center Göttingen, Göttingen, Germany. Email: [bjoern.tampe@med.uni-goettingen.de](mailto:bjoern.tampe@med.uni-goettingen.de)

Copyright © 2021 by the American Society of Nephrology

## FUNDING

None.

## ACKNOWLEDGMENTS

S. Hakroush and B. Tampe designed the study and wrote the original draft; S. Hakroush was responsible for the histopathological examination; and B. Tampe was responsible for the statistical analysis.

## DATA SHARING STATEMENT



The datasets used and/or analyzed during this study are available from the corresponding author on reasonable request.

## REFERENCES

1. Boudhabhay I, Delestre F, Coutance G, Gnemmi V, Quéméneur T, Vandenbussche C, et al.: Reappraisal of renal arteritis in ANCA-associated vasculitis: Clinical characteristics, pathology, and outcome. *J Am Soc Nephrol* 32: 2362–2374, 2021
2. Brix SR, Noriega M, Tennstedt P, Vettorazzi E, Busch M, Nitschke M, et al.: Development and validation of a renal risk score in ANCA-associated glomerulonephritis. *Kidney Int* 94: 1177–1188, 2018
3. Bonsib SM, Goeken JA, Fandel T, Houghton DC: Necrotizing medullary lesions in patients with ANCA associated renal disease. *Mod Pathol* 7: 181–185, 1994
4. Berden AE, Ferrario F, Hagen EC, Jayne DR, Jennette JC, Joh K, et al.: Histopathologic classification of ANCA-associated glomerulonephritis. *J Am Soc Nephrol* 21: 1628–1636, 2010

---

See related original article, "Reappraisal of Renal Arteritis in ANCA-associated Vasculitis: Clinical Characteristics, Pathology, and Outcome," in Vol. 32, Iss. 9, on pages 2362–2374.

Samy Hakroush <sup>1</sup> and Björn Tampe <sup>2</sup>

<sup>1</sup>Institute of Pathology, University Medical Center Göttingen, Germany

<sup>2</sup>Department of Nephrology and Rheumatology, University Medical Center Göttingen, Germany

*JASN* 2974–2975, 2021.

doi: <https://doi.org/10.1681/ASN.2021081057>